

STIC-ILL

RL 870.56

Pres

From: Holleran, Anne
Sent: Tuesday, September 04, 2001 6:03 PM
To: STIC-ILL
Subject: refs. for 09/251,133

Examiner: Anne Holleran
Art Unit: 1642; Rm 8E03
Phone: 308-8892
Date needed by: ASAP

36 2503

Please send me copies of the following :

1. Chien, J. et al. Mol. and Cell. Endocrinology (2001) 181(1-2): 69-79
2. Chien, J. et al. Int. J. of Cancer (2001) 91(1): 46-54
3. Chien, J. et al. Oncogene (1999) 18(22): 3376-3382
4. Wong, E.C.C. et al. Proc. Amer. Assoc. for Cancer Res. (1997) 38: 288
5. Rayford, W. et al. Prostate (1997) 30(3): 160-166
6. Xue-Zhang, Q. et al. Endocrine (1995) 3(6): 445-451
7. Shah, G.V. et al. Endocrinology (1994) 134(2): 596-602
8. Rayford, W. et al. J. of Urology (1994) 151(5 suppl): 490A
9. Rayford, W. et al. J. of Urology (1993) 149(4 suppl): 479A
10. Shah, G.V. et al. Prostate (N.Y.) (1992) 21(2): 87-97
11. Sagol, O. et al. Annals of Medical Sciences (1999) 8(1): 14-21
12. Sussenot, O. et al. Prostate (1998) 36(suppl. 8): 43-51
13. Hanna, F.W. et al. J. Endocrinol. (1997) 152(2): 275-281
14. Sim, S.J. et al. Annals of Clinical and Laboratory Science (1996) 26(6): 487-495
15. Watanabe, K. et al. Fukushima J. Medical Science (1995) 41(2): 141-152
16. Esik, O. et al. European J. Gynaecological Oncology (1994) 15(3): 211-216

JRL-100

WAW 9/21

The Journal of UROLOGY®

Official Journal of the American Urological Association, Inc.
Founded In 1917 By Hugh Hampton Young

**AUA EIGHTY-EIGHTH ANNUAL MEETING
MAY 15-20, 1993**

SAN ANTONIO CONVENTION CENTER - SAN ANTONIO, TEXAS

NOTICE: This material is protected
by copyright law (Title 17 U.S. Code)
Provided by the University of Washington Libraries



APR 08 1993

HEALTH SCIENCES LIBRARY

THE ALAMO

(Photo courtesy of the San Antonio Convention & Visitors Bureau)

Supplement to The Journal of Urology

BEST AVAILABLE COPY

TABLE OF CONTENTS

Officers	6A
Committees	
Executive Committee	6A
Local Arrangements Committee	7A
Ladies Hospitality Committee	7A
Program Committee	8A
Audio-Visual Committee	9A
Public Media Committee	9A
General Information	11A
Social Programs and Tours	15A
Corporate Patrons and Contributors	16A
Convention Floor Plan	18A
Exhibit Floor Plan	20A
Office of Education Courses	22A
Meeting Schedule	27A
Videotapes	36A
Videotape Library	43A
Urologic Topic Showcase Supporters	
Abbott Laboratories	51A
Hybritech, Inc.	51A
Ortho Pharmaceutical/McNeil Pharmaceutical Corporation	51A
SmithKline Beecham Pharmaceuticals	51A
Searle	52A
Wyeth-Ayerst	52A
American Medical Systems	120A
ZENECA Pharmaceuticals, A Business Unit of ZENECA, Inc. (Formerly ICI Pharma)	120A
Mentor Corporation	120A
Mission Pharmacal Company	120A
Osbon Medical Systems, Ltd.	120A
Merck Human Health Division	124A
Miles Inc., Pharmaceutical Division	124A
Ancillary Programs	
Society for Basic Urologic Research	44A
Urodynamics Society	45A
Society for Urology and Engineering	46A
International Scientific Committee on BPH	48A
Society of Urologic Oncology	49A
Society for the Study of Male Reproduction	50A
Society for Pediatric Urology (Podium Session 4)	59A
American Association of Clinical Urologists	132A
Scientific Program	
Sunday am	55A
Sunday pm	56A
Monday am	77A
Monday pm	78A
Tuesday am	99A
Tuesday pm	100A
Wednesday am	129A
Wednesday pm	143A
Thursday am	154A
Exhibits	164A
Abstracts	188A
Index of Participants	511A
Index of Authors	517A
Subject Index of Abstracts	544A

Important: The number preceding the title of a presentation is the abstract number.

ISSN 0022-5347

AUA Program

Copyright© 1993 American Urological Association, Inc.

(issued by Library of Congress)

Price \$25 per copy

1064

THE ARTIFICIAL NEURAL NETWORK: A NOVEL AND POWERFUL METHOD OF FERTILITY DATA ANALYSIS. C.S. Niederberger, L.I. Lipshultz, and D.J. Lamb, Houston, TX (Presentation to be made by Dr. Niederberger)

Analysis of fertility data is problematic for a variety of reasons: (1) although millions of sperm are ejaculated, the final step in conception requires only one functional sperm; (2) laboratory tests of fertility potential often reflect physiologic mechanisms that interact in complex ways; (3) patient data is often fragmented, and (4) 2 patients are required for the final result. Statistical methods in reproductive medicine are co-opted from other fields, such as oncology, and do not address the unique challenges of fertility data analysis. We are investigating a novel and powerful method of data analysis, the artificial neural network, in the assessment of fertility potential. The neural network is a computational algorithm inspired by the biological structure of the neuron, and "learns" to analyze data according to carefully chosen mathematical rules. To determine the validity of the neural network in analyzing tests of male reproductive function, we programmed a network using a common learning algorithm, backpropagation, to predict the results of sperm penetration assay (SPA) and penetration of sperm in bovine cervical mucus (Penetrak assay, Serono Labs) from the semen analysis (SA). The SPA and Penetrak tests are largely believed to be independent of the SA. Significantly, a neural network correctly predicted the Penetrak result in 80% of the assays that the network had not previously encountered, and another network predicted SPA outcome in nearly 70%. Results were compared to 2 traditional forms of statistical classification, linear and quadratic discriminant function analysis (LDFA and QDFA). The neural network was superior in predicting both assay outcomes. This classification algorithm may prove to serve many of the fundamental needs of data analysis unique to fertility studies.

1065

SKELETAL MUSCLE WRAP OF THE URINARY BLADDER AND FUNCTIONAL NEUROMUSCULAR ELECTRICAL STIMULATION: DETRUSOR-MYOPLASTY. Michael B. Chancellor¹, Rafael Acosta², Michael J. Erhard¹, David A. Rivas¹, Steven K. Salzman², Philadelphia, PA¹, Wilmington, DE². (Presentation to be made by Dr. Chancellor)

Detrusor-myoelastv (skeletal muscle assisted micturition) was investigated in a rat model of spinal cord injury. Anatomical dissection of the rectus muscle and bladder wrapping in 2 goats and 3 fresh human cadavers was also performed. The rectus muscle on one side of the abdomen was dissected free from the ventral fascia and transected cephalad above the level of the umbilicus. The muscular insertion to the symphysis pubis was left intact. Vascular flow via the inferior epigastric artery and vein was preserved. Innervation by 2 to 3 intercostal motor nerves was left intact.

Postoperatively, no bowel or abdominal wall functional deficit was apparent. The rotated muscular flap remained innervated and vascularized. No difference in 24 hr micturition patterns became evident between control rats and rats with a rectus muscle-wrapped bladder when not stimulated. Stimulation of the rectus muscle-wrapped bladder (both nerve and direct muscle stimulation) in the rat was capable of generating bladder pressure (range 10 to 60 cmH₂O) and achieved bladder emptying. Stimulation parameters ranged from 0.05-0.5 ms duration, 1-50 Hz, and 12.5-300 volts. Less voltage was required for nerve than muscle stimulation to achieve similar intravesical pressure. In both acute experiments and in rats surviving 1 month after spinal cord injury, sustained bladder contractions continued.

Dissection of goats and human cadavers revealed that a vascularized and innervated rectus muscle flap can be rotated into the pelvis and wrapped around the bladder without tension. This is the first report of the principle and technique of detrusor-myoelastv. Detrusor-myoelastv may be applicable for patients with an acontractile detrusor and non-intact sacral motor roots who are not candidates for sacral anterior root neurostimulation.

1066

EXPRESSION OF NEUROENDOCRINE FACTORS AND EXTRACELLULAR MATRIX DEGRADATIVE ENZYMES IN HUMAN PROSTATE TUMOR CELLS. N.M. Hoosein, C.J. Logothetis, M.G. Bandyk, G.L. Nicolson and L.W.K. Chung, Houston, TX. (presentation to be made by Dr. Hoosein).

Occurrence of neuroendocrine cells in prostate tumors has been associated with tumor progression (Abrahamsson et al., *Pathol. Res. Pract.* 185:373, 1989), early metastatic spread of the disease and poor prognosis (Tetu et al., *Cancer*, 59:1803, 1987; Cohen et al., *Br. J. Urol.*, 66:403, 1990). In this study, we have examined the presence of neuroendocrine substances chromogranin A (ChA), neuron-specific enolase (NSE), bombesin and serotonin in human prostate tumor cell lines and in bone marrow biopsies of prostate cancer patients. Previously, we had found that urokinase-type plasminogen activator (urokinase) plays an important role in determining the invasiveness of human prostate tumor cells (Hoosein et al., *Cancer Commun.* 3:255, 1991). Therefore, we also examined the presence of urokinase as well as another extracellular matrix degradative enzyme, heparanase, (Nakajima et al., *Science* 220:611, 1983) in prostate tumor cells. Results of immunocytochemical staining indicated that neuroendocrine markers ChA, NSE, and serotonin were expressed by prostate tumor cell lines of high metastatic potential, PC-3 and DU-145, but, not by the relatively indolent LNCaP cell line. High levels of bombesin-like immunoreactivity were found in PC-3 and DU-145 cells with lower levels in LNCaP cells. Also, PC-3 and DU-145 cells showed positive staining for urokinase and heparanase, whereas, LNCaP cells were negative for both these enzymes. Nests of tumor cells in bone marrow aspirates of two patients with prostatic adenocarcinoma, stained negative for bombesin, urokinase and heparanase, whilst, those in a patient with small cell carcinoma of the prostate were positive for all three markers. Our results indicate that PC-3 and DU-145 cells display neuroendocrine differentiation and that prostate tumor cells with neuroendocrine characteristics express high levels of degradative enzymes urokinase and heparanase, accounting for their aggressive behavior. Clinically, these neuroendocrine and enzyme markers may serve as valuable predictors of outcome and indicators for aggressive therapy.

1067

CALCITONIN RECEPTOR AFFECTS INTRACELLULAR CYCLIC 3', 5'-MONOPHOSPHATE (cAMP) LEVELS AND INCREASES CYTOPLASMIC Ca²⁺ CONCENTRATIONS IN HUMAN PROSTATE CANCER CELLS. *Walter Rayford, *Girish V. Shah, *Benjamin J. Frey, Mark J. Noble, Mark Austenfeld, John Weigcl, and Winston K. Mebust, Kansas City, KS (Presentation by Walter Rayford).

Presence of calcitonin (CT)-immunopositive cells have been identified in human prostate and its relative distribution increases in cancer. Our recent findings demonstrate that CT-like immunoreactive material (iCT) is secreted by primary prostate cells in culture, and its secretion from tumor cells is significantly higher than benign cells. Since CT and salmon CT (sCT)-like peptide act on a variety of organs through activation of different, and in some cases, multiple intracellular mechanisms, we have examined 1) whether CT receptors (CT-R) are present in human prostate tissue and human prostate cancer LNCaP cells; 2) the effect of CT on cAMP accumulation and/or cytoplasmic Ca²⁺ concentrations in LNCaP cells.

Surgical prostate specimens and LNCaP cells were homogenized to obtain plasma membrane fractions. The membranes were incubated with various concentrations of [¹²⁵I]-sCT to saturate the binding sites. The data were processed by Scatchard analysis. In a second group of experiments, cultured LNCaP cells were treated with various concentrations of sCT, and their intracellular cAMP levels were analyzed by RIA. In a third group of experiments, LNCaP cells were loaded with Fura 2AM and their intracellular Ca²⁺ levels were analyzed spectrofluorometrically.

Binding studies revealed presence of saturable CT binding sites in human prostate tissues and LNCaP cells. Binding affinity of CT-R in the tissue varied from 1 to 4.3 nM and the B_{max} ranged from 231-925 fmol/mg membrane protein. sCT demonstrated a biphasic effect on cAMP accumulation in LNCaP cells. At concentrations less than 1 nM, sCT significantly attenuated cAMP accumulation with maximal suppression of 30% at 10 pM. However, at concentrations greater than 1 nM, it significantly stimulated cAMP accumulation. The initial results suggest that at 100 nM, sCT significantly increased cytoplasmic Ca²⁺ levels in LNCaP cells. These findings suggest that CT-R is present in human prostate cells, and is coupled to adenylate cyclase and Ca²⁺ second messenger systems. When considered together with its secretion in the prostate, the results suggest a paracrine role for CT in pathophysiology of the prostate.